



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Question #5

What role can satellites take, as a complement to ground based measurement systems, to provide sustained observations to monitor GHG emissions?

Moustafa Chahine, Edward Olsen

**NOAA Hyperspectral
Spectrometer Workshop**

March 29 - 31, 2011

Miami Florida



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Question 5 – AIRS Specific

(concentrate on CO₂ here; Question 9 talk will cover other GHGs)

What role can satellites take, as a complement to ground based measurement systems, to provide sustained observations to monitor GHG emissions (e.g., CO₂, CH₄, O₃, N₂O, CFC's, NH₃, NF₃) that contribute to global warming

- **AIRS provides the increased spatial/temporal coverage required to identify regional carbon sources and sinks (especially in the Southern Hemisphere)**
Global coverage, day and night, cloudy and clear, all seasons
Provides integral constraint at coarse spatial scales and long time scales
- **AIRS partial column measurements complement ground based systems**
Mid-trop can be used to mitigate transport model errors (vertical and horizontal)
Extension to stratosphere will constrain CO₂ at the top end of the upward-looking FTIR averaging kernel and broaden understanding of Strat-Trop exchange
Extension to lower troposphere will enhance assimilation studies (esp. for SH)
- **What does AIRS show for future thermal IR atmosphere measurements?**
Higher spatial and spectral resolution required for lowest 1 km of atmosphere
Global coverage, day and night, cloudy and clear to lowest 1 km of atmosphere
- **The next GEN AIRS?**
ARIES concept for thermal sounder – expansion to shortwave for near surface₂



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

The Atmospheric Infrared Sounder on NASA's EOS Aqua Spacecraft


- AIRS Characteristics
- Launched: May 4, 2002
- Orbit: 705 km, 1:30pm, Sun Synch
- IFOV : $1.1^\circ \times 0.6^\circ$
(13.5 km x 7.4 km)
- Scan Range: $\pm 49.5^\circ$
- Full Aperture OBC Blackbody, $\varepsilon > 0.998$
- Full Aperture Space View
- Solid State Grating Spectrometer
 - IR Spectral Range:
3.74-4.61 μm , 6.2-8.22 μm ,
8.8-15.4 μm
 - IR Spectral Resolution:
 $\approx 1200 (\lambda/\Delta\lambda)$
 - # IR Channels: 2378 IR
- VIS Channels: 4
- Mass: 177Kg,
Power: 256 Watts,
Life: 5 years (7 years goal)
- Built: BAE Systems

AIRS



AIRS Spectra

AIRS Channels for Tropical Atmosphere with $T_{\text{surf}} = 301\text{K}$
Full Spectrum






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder

AIRS Products for Weather, Climate and Composition

AIRS Greenhouse Gases



Other AIRS Atmospheric Climate Products





National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

AIRS Operational Product Mid-Tropospheric CO₂ (8-10km)



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Global Yield of AIRS Level 2 Mid-Tropospheric CO₂


AIRS **Daily** CO₂ Yield 1°x1° Spatial Resolution

AIRS V5 CO2: Day 2003 7 15 x 1



AIRS **Monthly** CO₂ Yield 1°x1° Spatial Resolution

AIRS V5 CO2: Day 2003 7 15 x 30



AIRS Level 2 Mid-Tropospheric CO₂ retrieval yield is controlled by requirement for highest quality temperature and water vapor AIRS Level 2 products in 2x2 array of adjacent FOVs

A sounder with higher spatial and spectral resolution will increase yield and extend retrieved CO₂ profile to the near surface.

Thermal IR sounding allows retrievals day/night, pole-to-pole, land/ocean/ice, cloudy/clear




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Representative AIRS Mid-Trop CO₂ Averaging Kernels

(Individual AKs accompany each AIRS CO₂ sounding in the data products)






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

AIRS Data Show CO₂ is not well mixed in Mid-Troposphere

July 2003 AIRS mid trop CO₂ (5° smoothing) with 500 hPa gph contours



CO₂ is NOT Well Mixed in the mid-troposphere

- Driven by synoptic-scale phenomena (polar/subtropical jet streams)
- Complexity of the Southern Hemisphere not present in models
- AIRS mid-trop data will facilitate modeling of vertical & horizontal transport

M. Chahine et. al. (JPL)




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

AIRS - CarbonTracker upper-tropospheric CO₂ difference [ppm] vs. cloud top pressure eight N/S swaths, 2008



Davis Baker AGU 2010




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Comparison of Collocated AIRS CO₂ Retrievals with January 2009 HIPPO Data for profiles ranging from near surface to p < 200 hPa



HIPPO CO₂ vertical profile data courtesy of Steven C. Wofsy




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder


Collocated AIRS Mid-Trop CO₂ and Park Falls FTIR Total Column CO₂

AIRS and Park Falls Averaging Kernels



AIRS channels deliberately chosen
to suppress contribution from
surface and near-surface layers

$\Delta\text{CO}_2 = \langle \text{AIRS} \rangle - \text{Park Falls}$






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Seasonal Variation of Difference of Daily Average of Collocated AIRS Mid-Trop CO₂ and Park Falls FTIR



AIRS daytime data collated within radius of 500km of
highest quality Park Falls data taken within ± 2 hours of AIRS overpass




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder


Variability seen in 2009 HIPPO Campaign Compares well with AIRS Mid-Trop CO₂





S.C. Wofsy, et al (2011), HIAPER Pole-to-Pole Observations (HIPPO): Fine grained, global scale measurements of climatically important atmospheric gases and aerosols, *Proceedings of the Royal Society A*, in press.

AIRS has observed a
seasonally-variable
SH CO₂ Belt Since 2003


AIRS



GEOS Chem



M.T. Chahine, et al., Satellite remote sounding of mid-tropospheric CO₂, *Geophys. Res. Lett.*, 35, L17807, doi:10.1029/2008GL035022.



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Regional Comparison with CMDL surface observations

March – November 2005



Andrew Tangborn NASA/GSFC and UMBC

Assimilation of AIRS mid-trop CO₂ into GEOS5

Conclusions:

Comparison with CMDL surface data indicates that AIRS assimilation is improving the accuracy of surface values of CO₂ in GEOS5.

Differences between GEOS5 and AIRS CO₂ can be parameterized by hemisphere, with a systematic negative bias in the model during winter.



Observation Locations:

green = mean assimilation error is lower

Red = mean assimilation error is higher



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California



Atmospheric Infrared Sounder

Integrated Carbon Cycle Data Assimilation System

CAM3.5, LETKF Assimilating AIRS Mid-Trop CO₂



Junjie Liu UC Berkeley/JPL

Assimilation of AIRS mid-trop CO₂ improves spatial pattern




Spatial variability of AIRS-run is 0.53 ppm
which is much larger than that of the
Meteo-run (0.17 ppm)

Assimilation of AIRS mid-trop CO₂ adjusts vertical gradient May 2003: CO₂(850)-CO₂(400) (AIRS-run) - (Meteo-run)




NH: CO₂(850)>CO₂(400): fossil fuel+ land carbon source
SH: CO₂(850)<CO₂(400): transported from the NH
NOTE: scale of Met-run is 7x that of difference run


Assimilation of AIRS mid-trop CO₂ improves state estimate by 1 ppm



AIRS-run
assimilate CO₂
Met-run
no CO₂ obs



AIRS-run
assimilate CO₂
Met-run
no CO₂ obs



AIRS-run
assimilate CO₂
Met-run
no CO₂ obs






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Influences of El Niño on Mid-Trop CO₂ From AIRS and MOZART-2




Xun Jiang University of Houston



TOP: AIRS detrended and deseasonalized CO₂ anomaly averaged for 11 El Niño months

MIDDLE: AIRS detrended and deseasonalized CO₂ anomaly averaged for 17 La Niña months

BOTTOM: AIRS CO₂ anomaly difference (El Niño - La Niña)
(Consistent with change in Walker Circulation)



TOP: MOZART-2 CO₂ anomaly during El Niño

MIDDLE: MOZART-2 CO₂ anomaly during La Niña

BOTTOM: MOZART-2 CO₂ Difference (El Niño - La Niña)
(Signal is smaller than observed by AIRS)

Jiang, X., M. T. Chahine, E. T. Olsen, L. L. Chen, and Y. L. Yung (2010), Interannual variability of mid-tropospheric CO₂ from Atmospheric Infrared Sounder, Geophys. Res. Lett., 37, L13801, doi:10.1029/2010GL042823

NOTE: MOZART-2 results are preliminary. The boundary condition is a climatology and does not include interannual variability. (Courtesy of 16 Maochang Liang for the MOZART-2 model run)



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder



MJO-related AIRS Mid-Tropospheric CO₂ Anomaly Intraseasonal CO₂ variability across the global tropics

King-Fai Li, Tian, B., Waliser, D.E. and Yung, Y.L. (2010), Tropical mid-tropospheric CO₂ variability driven by the Madden-Julian Oscillation, PNAS, 107 (45), 19171-19175, doi: 10.1073/pnas.1008222107

MJO has previously been studied via its impact on atmospheric winds, pressure, temperature, moisture and rainfall.


Its impact upon mid-tropospheric CO₂ has now been detected.

This provides a new window of study of this planetary-scale zonal overturning circulation anomaly.



CO₂(ppmv)
-0.7 -0.6 -0.5 -0.4 -0.3 -0.2 -0.1 0.1 0.2 0.3 0.4 0.5 0.6 0.7
Contour lines: ECMWF-interim ω_{700} (Dotted: -ve, Solid: +ve)
Contours start from ± 4 mPa/s at an interval 4 mPa/s

The CO₂ anomaly is driven by the eastward-propagating vertical circulation of the MJO and implies that CO₂ values are higher at the surface than in the upper troposphere. This intraseasonal CO₂ variability provides a robustness test for chemical transport models.






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Mid-Trop CO₂ Bias due to H₂O Absorption is Small

- **Peak-to-Peak MJO Amplitude of H₂O at 600 hPa ≈ 1.4 g/kg**
[Tian et al. (2006), Vertical moist thermodynamic structure and spatial-temporal evolution of the MJO in AIRS observations, J. Atmos. Sci., 63, 2462]
- **Then Potential Bias in CO₂ $\approx 1.4 \times 0.13 < 0.2$ ppm**



King-Fai Li, Tian, B., Waliser, D.E. and Yung, Y.L. (2010), Tropical mid-tropospheric CO₂ variability driven by the Madden-Julian Oscillation, PNAS, 107 (45), 19171-19175, doi: 10.1073/pnas.1008222107



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Factors Affecting the CO₂ Retrievals

v range:	Mid-Troposphere -10km	Stratosphere – 30km	Lower Trop – 2.2km
	13 CO ₂ channels: 700 cm ⁻¹ – 722 cm ⁻¹	17 CO ₂ channels: 650 cm ⁻¹ – 680 cm ⁻¹	10 CO ₂ channels: 730 cm ⁻¹ – 745 cm ⁻¹
$T(p)$	Strong	Very strong	Strong
O ₃	Strong	Weak	Medium
H ₂ O	Medium	No impact	Medium
Surface emission, E _s (T _s , ε _s)	Very weak	No impact	Medium
$\Delta G/\Delta \text{CO}_2^*$	~0.4	~0.2	~0.5

*($\Delta G/\Delta \text{CO}_2$) describes the sensitivity of observed spectra to changes in CO₂. It is a function of the lapse rate of atmospheric temperature profiles which is 7 K/km in the mid-troposphere, 1.5K/km in the stratosphere and 10K/km near surface.



- **Mid-troposphere: Operational and Released to the Public (Sept 2002 – Present)**
- **Stratosphere: Algorithm Completed, QA and Validation Underway (8/2010)**
- **Lower troposphere: Algorithm Nearly Complete, Preliminary Retrievals Underway (12/2010)**




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

3 Layers of CO₂ Derived from AIRS


Summary




Stratosphere

Mid-Troposphere


Lower Trop



Preliminary



Complete
Sept 02 - Present



Preliminary



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

3 Layers of CO₂ Derived from AIRS

Current Status

Task	Mid-Troposphere	Stratosphere	Lower Troposphere
Algorithm Development Initial Channel Selection	✓	✓	✓
Retrieval Optimization Beta Software Development & Test Refine Channel Selection Refine Quality Control	✓	✓	✓
Validation and Comparison In-Situ Measurements Models	✓*	In progress	In progress
Report Results Professional Meetings Journal Publications	✓		
Transition to Operational Stage Production Software Development Documentation	✓		
Production	✓		
Distribution via GES DISC & JPL	✓*		

*Continuous Updates

M. Chahine et. al. (JPL)



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

ARIES can map GHG emissions from large cities and counties

ARIES Characteristics:



- Extension of AIRS Methodology
- Global Maps Daily ($\pm 55^\circ$ Swath)
- Spatial Resolution: 2 km
- Spectral Range: 3.0 – 15.4 μm
- Spectral Resolution: 0.5 cm^{-1}

Over 5000 Channels



Products:

Vertical Profiles of:

T, H₂O, CO₂, CH₄, CO, N₂O, SO₂,
HNO₃, O₃



AIRS CO₂ Map, July 2003



ARIES CO₂ Map Resolution

Products	IFOV (km)	λ_1 (μm) ν_1 (cm^{-1})	λ_2 (μm) ν_2 (cm^{-1})	R, $\Delta\nu$ (cm^{-1})
Temperature, CO ₂ , CH ₄ , N ₂ O, CO	1	3.39 2950	4.76 2100	2.0
Water, CH ₄ , SO ₂ , HNO ₃	1	6.20 1613	8.70 1150	1.0
O ₃ , HNO ₃	1	8.70 1150	11.36 880	0.5
Temperature, CO ₂	1	11.36 880	15.38 650	0.5

T. Pagano et. al. (JPL)



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Moustafa T. Chahine

1935 - 2011

